Abstract Submitted for the 1995 Meeting of the American Physical Society San Jose, California March 20-24, 1995

Suggested Session Title:Optical and Magnetic Properties of YBCORelated Structures

Classification Scheme:

12g

Raman Study of Pr doped Eu_{1.5}Ce_{0.5}Sr₂Cu₂MO₁₀ (M = Nb and Ta) High T_C Superconductors,* -- M. J. BENNAHMIAS, 1,2 H. B. RADOUSKY, 2,3 H. E. LORENZANA, 2 T.J. GOODWIN,³ AND R.N. SHELTON,³ ¹UC-Davis, Dept. of Applied Science, Livermore, CA 94551, ²Lawrence Livermore National Laboratory, Livermore, CA 94551, ³UC-Davis, Dept. of Physics, Davis, Ca 95616 -- Pr_{1.5}Ce_{0.5}Sr₂Cu₂MO₁₀ (PrCeSCMO) (M = Nb and Ta) are insulating analogs to the family of cuprate-niobate superconducting compounds RCeSCNO (R = Eu, Sm, and Nd) with TC's near 28 K.¹ Raman characterization has been performed on a wide range of $(Eu_{1.5-x}Pr_x)CeSCMO$ (M = Nb and Ta; 0 < x < 1.5) compounds. The phase diagram for these compounds indicates that for x < 0.3 the compounds are superconducting and for x > 0.6 complex magnetic behavior is observed. The line assignments for these Raman spectra are similar to the parent compound RBa₂Cu₃O₇. The origin of the broadening of the Raman peaks upon Pr doping will be discussed in terms of disorder within the CuO planes. The effect of this disorder on promoting the weak ferromagnetism seen in these materials will also be briefly discussed.

*Work at LLNL performed under the auspices of the USDoE under contract number W-7405-ENG-48. Work at UC Davis was partially funded under NSF grant no. DMR94-03895.

[1]. T.J. Goodwin, H. B. Radousky and R. N. Shelton, Physica C 204, 212 (1992).